

VOLATILE COMPOUNDS RESPONSIBLE FOR THE FISHY OFF-FLAVOUR IN BEEF WITH ENHANCED HEALTHFUL FATTY ACIDS

N. Prieto*, R. Thacker, I.L. Larsen, M.E.R. Dugan, P. Vahmani, M. Juárez, Ó. López-Campos
and J.L. Aalhus

Agriculture and Agri-Food Canada, Lacombe, AB, T4L 1W1, Canada

*Corresponding author email: nuria.prietobenavides@canada.ca

I. INTRODUCTION

There have been many attempts to increase n-3 fatty acids (FA) in beef due to their health benefits and their presence in ruminant animals [1]. However, when polyunsaturated fatty acids are too abundant, oxidation can result in a shorter shelf life and negative flavours or off-flavours in cooked meat. Vahmani et al. [2] improved the profile of health favourable FA (n-3, vaccenic and rumenic) in beef by feeding co-extruded flaxseed (flaxseed, peas, and alfalfa) before alfalfa-grass hay. However, this beef enriched with healthful FA resulted in tougher steaks with lower beef flavour and greater off-flavour intensity, presenting a fishy off-flavour to an extent that might be detectable by consumers [3]. The aim of this study was to characterize the volatile compounds responsible for the fishy off-flavour found in beef with health enhanced FA profiles.

II. MATERIALS AND METHODS

Twelve 2.5-cm steaks enriched with healthful FA presenting a fishy off-flavour according to trained panellists (Fishy) [3], and 11 steaks from commercial cattle (Non-fishy) collected from the caudal end of the *longissimus thoracis* were frozen at -20 °C. Upon thawing at 4 °C for 24 h, steaks were cooked to a final internal temperature of 71 °C [3]. Stir bar sorptive extraction (SBSE) coupled with thermal desorption-gas chromatography-mass spectrometry (TD-GC-MS) was used for identification and quantification of volatile flavour compounds in grilled beef samples as described in Ruan et al. [4]. Partial least squares discriminant analysis (PLS-DA) based on volatile compounds was applied to classify steaks into presence or absence of fishy off-flavour (Fishy and Non-fishy, respectively).

III. RESULTS AND DISCUSSION

The PLS-DA including only two factors identified two clusters: the first group included Fishy samples and the second cluster Non-fishy samples. Hence, a clear separation between Fishy and Non-fishy was observed based on the volatile compounds.





Figure 1. Score plot based on factor 1 and 2 from the partial least squares discriminant analysis using the volatile compounds to classify steaks based on the presence or absence of fishy off-flavour.

As shown in Fig. 2, 17 volatile compounds significantly ($P < 0.05$) contributed to the presence of fishy off-flavour. Among these, 14 compounds were positioned on the right quadrants near the fishy off-flavour, hence being positively correlated with the presence of this off-flavour. Compounds produced by autoxidation of linolenic acid, such as (E,E)-2,4-Heptadienal and heptanal, and 2-undecanone and pentadecane have been described as major contributors to fishy odour [5,6]. To our knowledge, the other 10 compounds have not been independently related to fishy off-flavour, although they might have



contributed to this off-flavour when their odour was mixed with others in a complex beef sample.

Figure 2. Correlation loading plot from the partial least squares discriminant analysis showing the volatile compounds positively and negatively correlated with the presence of fishy off-flavour. Circled (•) volatile compounds were significant at $P < 0.05$.

IV. CONCLUSION

A variety of compounds appears to have contributed towards the fishy off-flavour found in beef with enhanced healthful FA. Fishy off-flavour is a complex sensory phenomenon, and further research to confirm the contribution of those volatile compounds not previously related to this off-flavour is warranted.

ACKNOWLEDGEMENTS

This project was funded by the Alberta Livestock and Meat Agency (2015P006R). The gift of LinPro™-R from O&T Farms (Regina, SK, Canada) is gratefully acknowledged. The authors also thank the Lacombe Research and Development Centre operational, processing and technical staff for their dedication and expert assistance.

REFERENCES

1. Field, C. J., Blewett, H. H., Proctor, S., & Vine, D. (2009). Human health benefits of vaccenic acid. *Applied Physiology, Nutrition, and Metabolism* 34(5), 979-991.
2. Vahmani, P., Rolland, D. C., McAllister, T. A., Block, H. C., Proctor, S. D., Guand, L. L., Prieto, N., Aalhus, J. L., & M. E. R. Dugan (2017). Effects of feeding steers extruded flaxseed mixed with hay or before hay on animal performance, carcass quality, and meat and hamburger fatty acid composition. *Meat Science* 131, 9-17.
3. Prieto, N., Dugan, M. E. R., Larsen, I. L., Vahmani, P., & Aalhus, J. L. (2017). Palatability of beef from cattle fed extruded flaxseed before hay or mixed with hay. *Meat and Muscle Biology* 1, 157-168.

4. Ruan, E. D., Aalhus, J. L., Juarez, M., & Sabik, H. (2015). Analysis of volatile and flavor compounds in grilled lean beef by stir bar sorptive extraction and thermal desorption-gas chromatography Mass Spectrometry. *Food Analytical Methods* 8, 363-370.
5. Giri, A., Osako, K., & Ohshima T. (2010). Identification and characterisation of headspace volatiles of fish miso, a Japanese fish meat based fermented paste, with special emphasis on effect of fish species and meat washing. *Food Chemistry* 120, 621-631.
6. Crawford, L. & Kretsch, M. J. (1976). GC-MS identification of the volatile compounds extracted from roasted turkeys fed a basal diet supplemented with tuna oil: some comments on fishy flavor. *Journal of Food Science* 41(6), 1470-1478.